

# $C^*$ -algebras generated by radial Toeplitz operators on the Bergman and Fock spaces

Egor Maximenko


based on joint works with Kevin Esmeral, Sergei Grudsky,  
Crispin Herrera Yañez and Nikolai Vasilevski

Instituto Politécnico Nacional, ESFM, México

Taller “Análisis: Norte - Sur '16”  
CINVESTAV del IPN, México  
16–18 de noviembre de 2016

Oscillating sequences

Fock case



Approximation

Bergman case

# Bounded uniformly continuous functions

## Modulus of continuity

Given  $f: \mathbb{R} \rightarrow \mathbb{C}$ , define  $\omega_f: [0, +\infty) \rightarrow [0, +\infty)$  by

$$\omega_f(\delta) := \sup\{|f(x) - f(y)|: x, y \in \mathbb{R}, |x - y| \leq \delta\}.$$

## Bounded uniformly continuous functions

$$C_u(\mathbb{R}) := \left\{ f \in C^{\mathbb{R}}: \sup_{x \in \mathbb{R}} |f(x)| < +\infty \quad \wedge \quad \lim_{\delta \rightarrow 0^+} \omega_f(\delta) = 0 \right\}.$$

**Proposition.**  $C_u(\mathbb{R})$  is a  $C^*$ -subalgebra of  $L^\infty(\mathbb{R})$ .

# Convolution, Fourier transform

## Convolution

Given  $k \in L^1(\mathbb{R})$  and  $f \in L^\infty(\mathbb{R})$ ,

$$(k * f)(x) := \int_{\mathbb{R}} k(x - y)f(y) dy.$$


It is easy to see that  $k * f \in C_u(\mathbb{R})$ .

## Fourier transform

Given  $k \in L^1(\mathbb{R})$ ,

$$\widehat{k}(t) := \int_{\mathbb{R}} k(x) e^{-2\pi i xt} dx.$$

## Convolution and oscillations: example



## Bounded uniformly continuous functions can be uniformly approximated by convolutions

**Approximation Theorem.** Let  $k \in L^1(\mathbb{R})$  satisfy Wiener's condition:

$$\forall t \in \mathbb{R} \quad \widehat{k}(t) \neq 0.$$


Then  $\{k * f : f \in L^\infty(\mathbb{R})\}$  is a dense subset of  $C_u(\mathbb{R})$ .

We have not found this result in the literature, so we included a proof in Esmeral, Maximenko (2016) doi:10.1007/s11785-016-0557-0

The theorem is very close to Tauberian Wiener's theorem, and our proof is based on the well-known "Wiener's approximate deconvolution technique": Dirac sequences and Wiener's Division Lemma.

Oscillating sequences

Fock case



Approximation

Bergman case

## Log-oscillating sequences (Robert Schmidt, 1924)

**Definition.** LO := bounded sequences  $x = (x_n)_{n=0}^\infty$  such that

$$\lim_{\substack{m+1 \\ n+1} \rightarrow 1} |x_m - x_n| = 0.$$

In other words, LO is the set of all bounded functions  $\mathbb{N}_0 \rightarrow \mathbb{C}$  that are uniformly continuous with respect to the log-metric:

$$\rho_{\log}(m, n) := |\log(m+1) - \log(n+1)|,$$

$$\omega_{\rho_{\log}, x}(\delta) := \sup\{|x_m - x_n| : m, n \in \mathbb{N}_0, \rho_{\log}(m, n) \leq \delta\},$$

$$\text{LO} := \{x \in \ell^\infty : \lim_{\delta \rightarrow 0^+} \omega_{\rho_{\log}, x}(\delta) = 0\}.$$

**Proposition.** LO is a  $C^*$ -subalgebra of  $\ell^\infty$ .



## Log-oscillating sequences can be obtained from bounded uniformly continuous functions

**Proposition.** The operator  $\Phi: C_u(\mathbb{R}) \rightarrow \text{LO}$ , defined by

$$(\Phi f)_n := f(\log(n+1)),$$

is an epimorphism of  $C^*$ -algebras.

**Idea to prove the surjective property:**

Given a sequence  $x \in \text{LO}$ , put

$$f(\log(n+1)) := x_n \quad (n \in \mathbb{N}_0)$$

and apply the linear interpolation between the nodes  $\log(n+1)$ .

## Sqrt-oscillating sequences

RO := the set of all bounded sequences that are uniformly continuous with respect to the sqrt-metric:

$$\rho_{\text{sqrt}}(m, n) := |\sqrt{m} - \sqrt{n}|.$$

**Proposition.** RO is a  $C^*$ -subalgebra of  $\ell^\infty$ .


**Proposition.** The operator  $\Psi: C_u(\mathbb{R}) \rightarrow \text{RO}$ , defined by

$$(\Psi f)_n := f(\sqrt{n}),$$


is an epimorphism of  $C^*$ -algebras.

## Example of log-oscillation and sqrt-oscillation

$$x_n = \cos(\log(n + 1)):$$




$$x_n = \cos(\sqrt{n}):$$



Oscillating sequences

Fock case



Approximation

Bergman case

## Bergman space on the unit disk

$$\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}.$$

$$\mu_{\mathbb{D}} := \frac{1}{\pi} \text{ Lebesgue plane measure.}$$

$$\mathcal{A}^2(\mathbb{D}) := \{f \in H(\mathbb{D}) : f \in L^2(\mathbb{D})\}.$$

Bergman projection  $P_{\mathbb{D}} :=$  orthogonal projection  $L^2(\mathbb{D}) \rightarrow \mathcal{A}^2(\mathbb{D})$ .

Orthonormal basis in  $\mathcal{A}^2(\mathbb{D})$  consisting of normalized monomials:

$$e_n(z) = \sqrt{n+1} z^n.$$

Given  $g \in L^\infty(\mathbb{D})$ , denote by  $T_g$  the Toeplitz operator

$$T_g : \mathcal{A}^2(\mathbb{D}) \rightarrow \mathcal{A}^2(\mathbb{D}), \quad T_g f := P_{\mathbb{D}}(gf).$$

## Radial Toeplitz operators on the Bergman space

Given  $a \in L^\infty([0, 1])$ , denote by  $\tilde{a}$  its extension onto  $\mathbb{D}$ :

$$g(z) := \tilde{a}(z) := a(|z|) \quad (z \in \mathbb{D}).$$

It is easy to see that Toeplitz operators generated by radial symbols are diagonal in the monomial basis:

**Diagonalization of radial Toeplitz operators on the Bergman space:**

$$T_g e_n = \lambda_a(n) e_n,$$

where

$$\lambda_a(n) = (n + 1) \int_0^1 a(\sqrt{r}) r^n dr.$$

Korenblum, Zhu (1995)

<http://www.mathjournals.org/jot/1995-033-002/1995-033-002-010.html>

# From radial Toeplitz operators to eigenvalues' sequences

$$\begin{array}{ccc} T_{\tilde{a}} & \longleftrightarrow & \lambda_a \\ \{T_{\tilde{a}}: a \in L^\infty([0,1])\} & \longleftrightarrow & \Lambda := \{\lambda_a: a \in L^\infty([0,1])\} \\ C^*\text{-alg}\{T_{\tilde{a}}: a \in L^\infty(0,1)\} & \longleftrightarrow & C^*\text{-alg}(\Lambda) \end{array}$$

## Natural questions:

What is the closure of  $\Lambda$  ?

What is the  $C^*$ -algebra generated by  $\Lambda$  ?

# From radial Toeplitz operators to eigenvalues' sequences

$$\begin{array}{ccc} T_{\tilde{a}} & \longleftrightarrow & \lambda_a \\ \{T_{\tilde{a}}: a \in L^\infty([0, 1))\} & \longleftrightarrow & \Lambda := \{\lambda_a: a \in L^\infty([0, 1))\} \\ C^* \text{-alg}\{T_{\tilde{a}}: a \in L^\infty(0, 1)\} & \longleftrightarrow & C^* \text{-alg}(\Lambda) \end{array}$$

## Natural questions:

What is the closure of  $\Lambda$  ?

What is the  $C^*$ -algebra generated by  $\Lambda$  ?

These two questions have the same answer: LO.



## Key idea: spectral sequences as convolutions

$$\lambda_a(n) = (n+1) \int_0^1 a(\sqrt{r}) r^n dr.$$

Change of variables:

$$n+1 = \exp(x), \quad r = \exp(-\exp(-y)).$$


$$\lambda_a(n) = \int_{\mathbb{R}} \underbrace{a(\exp(-\exp(-y)/2))}_{b(y)} k(x-y) dy = (k * b)(x),$$

where

$$k(x) = \frac{e^x}{e^{e^x}}.$$

## The convolution kernel satisfies Wiener's condition

$$k(x) = \frac{e^x}{e^{e^x}}, \quad \widehat{k}(t) = \Gamma(1 - 2i\pi t) \neq 0 \quad (t \in \mathbb{R}).$$



## Main result for the radial Bergman case

**Theorem.**  $\underbrace{\{\lambda_a: a \in L^\infty([0, 1])\}}_{\Lambda}$  is a dense subset of LO.

As a consequence,  $C^*(\Lambda) = \text{LO}$ , and the  $C^*$ -algebra generated by radial Toeplitz operators on  $\mathcal{A}^2(\mathbb{D})$  is isometrically isomorphic to LO.

Suárez (2008) doi:10.1112/blms/bdn042


Grudsky, M, Vasilevski (2013) euclid.cma/1356039033

Bauer, Herrera Yañez, Vasilevski (2014) doi:10.1007/s00020-013-2101-1


Herrera Yañez, M, Vasilevski (2015) doi:10.1007/s00020-014-2213-2

## Example of spectral sequence in radial Bergman case

The function  $a(r) = \cos(\log(-2 \log(r)))$  oscillates rapidly near  $r = 1$ .




Then  $\lambda_a(n) = \operatorname{Re}(\Gamma(1 - i) \exp(i \log(n + 1)))$ .



Oscillating sequences

Fock case



Approximation

Bergman case

## Bargmann–Segal–Fock space

Gaussian measure  $d\gamma(z) = \frac{1}{\pi} e^{-|z|^2} d\mu(z)$ .

$\mathcal{A}^2(\mathbb{C}, d\gamma) := \{f \in H(\mathbb{C}) : f \in L^2(\mathbb{C}, d\gamma)\}$ .

$P_{\mathbb{C}} :=$  orthogonal projection  $L^2(\mathbb{C}, d\gamma) \rightarrow \mathcal{A}^2(\mathbb{C}, d\gamma)$ .

Orthonormal basis in  $\mathcal{A}^2(\mathbb{C}, d\gamma)$ :

$$f_n(z) = \frac{z^n}{\sqrt{n!}}.$$

Given  $g \in L^\infty(\mathbb{C})$ , denote by  $T_g$  the Toeplitz operator

$$T_g : \mathcal{A}^2(\mathbb{C}, d\gamma) \rightarrow \mathcal{A}^2(\mathbb{C}, d\gamma), \quad T_g f := P_{\mathbb{C}}(gf).$$

# Radial Toeplitz operators on the Fock space

Given  $a \in L^\infty(\mathbb{R}_+)$ , denote by  $\tilde{a}$  its extension onto  $\mathbb{C}$ :

$$g(z) := \tilde{a}(z) := a(|z|) \quad (z \in \mathbb{C}).$$

**Diagonalization of radial Toeplitz operators on the Fock space:**

$$T_g f_n = \xi_a(n) f_n,$$

where

$$\xi_a(n) = \frac{1}{\sqrt{n!}} \int_{\mathbb{R}_+} a(\sqrt{r}) e^{-r} r^n dr.$$

Grudsky, Vasilevski (2002) doi:10.1007/BF01197858

## Key idea: approximate spectral sequences by convolutions

We proved that

$$\xi_a(n) - \sqrt{\frac{2}{\pi}} \int_{\mathbb{R}_+} e^{-2(\sqrt{n}-y)^2} a(y) dy \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Let  $a_{\text{ext}}$  be the function  $a$  extended from  $\mathbb{R}_+$  onto  $\mathbb{R}$  by zero. Then

$$\xi_a(n) \approx (k * a_{\text{ext}})(\sqrt{n}) \quad \text{as } n \rightarrow \infty,$$

where  $k$  is the following Gaussian density:

$$k(x) = \sqrt{\frac{2}{\pi}} e^{-2x^2}.$$

The Fourier transform of  $k$  does not vanish:

$$\widehat{k}(t) = e^{-\frac{1}{2}\pi^2 t^2}.$$



## Main result for the radial Fock case

**Theorem.**  $\{\xi_a: a \in L^\infty(\mathbb{R}_+)\}$  is a dense subset of RO.

As a consequence, the  $C^*$ -algebra generated by radial Toeplitz operators on  $\mathcal{A}^2(\mathbb{C}, d\gamma)$  is isometrically isomorphic to RO.


Esmeral, Maximenko (2016) doi:10.1007/s11785-016-0557-0

## Example of spectral sequence in radial Fock case

Let  $a(r) = \cos(r)$ ,  $r \in \mathbb{R}_+$ .

Then the eigenvalue sequence  $\xi_a$  has a typical  $\sqrt{\cdot}$ -oscillation:

$$\xi_a(n) = {}_1F_1(1+n; 1/2; 1/4) \approx e^{-1/8} \cos(\sqrt{n}) + o(1).$$



# Summary

